

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
21 November 2002 (21.11.2002)

PCT

(10) International Publication Number
WO 02/092083 A1

(51) International Patent Classification⁷: **A61K 31/4188**,
31/4196, 31/424, 31/429, 31/436, 31/4365, 31/4375,
31/4985, 31/5365, C07D 471/04, 491/052, 487/04

(21) International Application Number: PCT/US02/15398

(22) International Filing Date: 16 May 2002 (16.05.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/290,976 16 May 2001 (16.05.2001) US

(71) Applicant: **CYTOVIA, INC.** [US/US]; 6650 Nancy
Ridge Drive, San Diego, CA 92121 (US).

(71) Applicants and

(72) Inventors: **CAI, Sui, Xiong** [US/US]; 3623 Berry-
field Court, Sand Diego, CA 92130 (US). **XU, Lifan**
[CN/US]; 10117 Caminito Zar, San Diego, CA 92126
(US). **STORER, Richard** [GB/GB]; 26 Cecil Park,
Pinner, Middlesex HA5 5HH (GB). **ATTARDO, Giorgio**
[CA/CA]; 2740 Prudential, Laval, Quebec H7K 3M1
(CA).

(74) Agents: **ESMOND, Robert, W. et al.**; Sterne, Kessler,
Goldstein & Fox P.L.L.C., 1100 New York Avenue, N.W.,
Suite 600, Washington, DC 20005-3934 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN,
YU, ZA, ZM, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: 7,8-FUSED 4S(H)-CHROMENE AND ANALOGS AS ACTIVATORS OF CASPASES AND INDUCERS OF APOPTOSIS AND THE USE THEREOF



(I)

(57) Abstract: The present invention is directed to substituted 4H-chromene and analogs thereof, represented by the general formula (I) or pharmaceutically acceptable salts or prodrugs thereof, wherein: A, R₁, R₂, R₅, X, Y, and Z, are defined herein and B is fused thiazole, oxazole, 2-imino-imidazole, 2,1,3-thiadiazol-2-one, thiazol-2-one, oxazol-2-one, imidazol-2-thione, thiazol-2-thione, oxazol-2-thione, imidazoline, oxazoline, thiazoline, triazole, oxazine, oxazine-2,3-dione, or piperazine ring. The present invention also relates to the discovery that compounds having Formula I are activators of caspases and

inducers of apoptosis. Therefore, the activators of caspases and inducers of apoptosis of this invention can be used to induce cell death in a variety of clinical conditions in which uncontrolled growth and spread of abnormal cells occurs.

WO 02/092083 A1

7,8-FUSED 4H-CHROMENE AND ANALOGS AS ACTIVATORS OF
CASPASES AND INDUCERS OF APOPTOSIS AND THE USE THEREOF

BACKGROUND OF THE INVENTION

Field of the Invention

5 This invention is in the field of medicinal chemistry. In particular, the invention relates to 7,8-fused 4H-chromene and analogs, and the discovery that these compounds are activators of caspases and inducers of apoptosis. The invention also relates to the use of these compounds as therapeutically effective anti-cancer agents.

10 Related Art

 Organisms eliminate unwanted cells by a process variously known as regulated cell death, programmed cell death, or apoptosis. Such cell death occurs as a normal aspect of animal development as well as in tissue homeostasis and aging (Glucksmann, A., *Biol. Rev. Cambridge Philos. Soc.* 26:59-86 (1951); Glucksmann, A., *Archives de Biologie* 76:419-437 (1965);
15 Ellis, *et al.*, *Dev.* 112:591-603 (1991); Vaux, *et al.*, *Cell* 76:777-779 (1994)). Apoptosis regulates cell number, facilitates morphogenesis, removes harmful or otherwise abnormal cells and eliminates cells that have already performed their function. Additionally, apoptosis occurs in response to various
20 physiological stresses, such as hypoxia or ischemia (PCT published application WO96/20721).

 There are a number of morphological changes shared by cells experiencing regulated cell death, including plasma and nuclear membrane blebbing, cell shrinkage (condensation of nucleoplasm and cytoplasm),
25 organelle relocation and compaction, chromatin condensation and production of apoptotic bodies (membrane enclosed particles containing intracellular material) (Orrenius, S., *J. Internal Medicine* 237:529-536 (1995)).

 Apoptosis is achieved through an endogenous mechanism of cellular suicide (Wyllie, A.H., in *Cell Death in Biology and Pathology*, Bowen and

Lockshin, eds., Chapman and Hall (1981), pp. 9-34). A cell activates its internally encoded suicide program as a result of either internal or external signals. The suicide program is executed through the activation of a carefully regulated genetic program (Wyllie, *et al.*, *Int. Rev. Cyt.* 68:251 (1980); Ellis, *et al.*, *Ann. Rev. Cell Bio.* 7:663 (1991)). Apoptotic cells and bodies are usually recognized and cleared by neighboring cells or macrophages before lysis. Because of this clearance mechanism, inflammation is not induced despite the clearance of great numbers of cells (Orrenius, S., *J. Internal Medicine* 237:529-536 (1995)).

It has been found that a group of proteases are a key element in apoptosis (see, e.g., Thornberry, *Chemistry and Biology* 5:R97-R103 (1998); Thornberry, *British Med. Bull.* 53:478-490 (1996)). Genetic studies in the nematode *Caenorhabditis elegans* revealed that apoptotic cell death involves at least 14 genes, 2 of which are the pro-apoptotic (death-promoting) *ced* (for *cell death abnormal*) genes, *ced-3* and *ced-4*. CED-3 is homologous to interleukin 1 beta-converting enzyme, a cysteine protease, which is now called caspase-1. When these data were ultimately applied to mammals, and upon further extensive investigation, it was found that the mammalian apoptosis system appears to involve a cascade of caspases, or a system that behaves like a cascade of caspases. At present, the caspase family of cysteine proteases comprises 14 different members, and more may be discovered in the future. All known caspases are synthesized as zymogens that require cleavage at an aspartyl residue prior to forming the active enzyme. Thus, caspases are capable of activating other caspases, in the manner of an amplifying cascade.


Apoptosis and caspases are thought to be crucial in the development of cancer (*Apoptosis and Cancer Chemotherapy*, Hickman and Dive, eds., Humana Press (1999)). There is mounting evidence that cancer cells, while containing caspases, lack parts of the molecular machinery that activates the caspase cascade. This makes the cancer cells lose their capacity to undergo cellular suicide and the cells become immortal and cancerous. In the case of the apoptosis process, control points are known to exist that represent points for intervention leading to activation. These control points include the CED-9-BCL-like and CED-3-ICE-like gene family products, which are intrinsic

proteins regulating the decision of a cell to survive or die and executing part of the cell death process itself, respectively (see, Schmitt, *et al.*, *Biochem. Cell. Biol.* 75:301-314 (1997)). BCL-like proteins include BCL-xL and BAX-alpha, which appear to function upstream of caspase activation. BCL-xL appears to prevent activation of the apoptotic protease cascade, whereas BAX-alpha accelerates activation of the apoptotic protease cascade.

It has been shown that chemotherapeutic (anti-cancer) drugs can trigger cancer cells to undergo suicide by activating the dormant caspase cascade. This may be a crucial aspect of the mode of action of most, if not all, known anticancer drugs (Los, *et al.*, *Blood* 90:3118-3129 (1997); Friesen, *et al.*, *Nat. Med.* 2:574 (1996)). The mechanism of action of current antineoplastic drugs frequently involves an attack at specific phases of the cell cycle. In brief, the cell cycle refers to the stages through which cells normally progress during their lifetime. Normally, cells exist in a resting phase termed G₀. During multiplication, cells progress to a stage in which DNA synthesis occurs, termed S. Later, cell division, or mitosis occurs, in a phase called M. Antineoplastic drugs, such as cytosine arabinoside, hydroxyurea, 6-mercaptopurine, and methotrexate are S phase specific, whereas antineoplastic drugs, such as vincristine, vinblastine, and paclitaxel are M phase specific. Many slow growing tumors, e.g., colon cancers, exist primarily in the G₀ phase, whereas rapidly proliferating normal tissues, e.g., bone marrow, exist primarily in the S or M phase. Thus, a drug like 6-mercaptopurine can cause bone marrow toxicity while remaining ineffective for a slow growing tumor. Further aspects of the chemotherapy of neoplastic diseases are known to those skilled in the art (see, e.g., Hardman, *et al.*, eds., *Goodman and Gilman's The Pharmacological Basis of Therapeutics*, Ninth Edition, McGraw-Hill, New York (1996), pp. 1225-1287). Thus, it is clear that the possibility exists for the activation of the caspase cascade, although the exact mechanisms for doing so are not clear at this point. It is equally clear that insufficient activity of the caspase cascade and consequent apoptotic events are implicated in various types of cancer. The development of caspase cascade activators and inducers of apoptosis is a highly desirable goal in the development of therapeutically effective antineoplastic agents. Moreover,

since autoimmune disease and certain degenerative diseases also involve the proliferation of abnormal cells, therapeutic treatment for these diseases could also involve the enhancement of the apoptotic process through the administration of appropriate caspase cascade activators and inducers of apoptosis.

EP537949 discloses derivatives of 4*H*-naphtho[1,2-*b*]pyran as antiproliferatives:




wherein,

each R^1 is independently halo, trifluoromethyl, C_{1-4} alkoxy, hydroxy, nitro, C_{1-4} alkyl, C_{1-4} alkylthio, hydroxy- C_{1-4} alkyl, hydroxy- C_{1-4} alkoxy, trifluoromethoxy, carboxy, $-COOR^5$ where R^5 is an ester group, $-CONR^6R^7$ or $-NR^6R^7$ where R^6 and R^7 are each hydrogen or C_{1-4} alkyl;

R^2 is phenyl, naphthyl or heteroaryl selected from thienyl, pyridyl, benzothienyl, quinoliny, benzofuranyl or benzimidazolyl, wherein said phenyl, naphthyl and heteroaryl groups are optionally substituted, or R^2 is furanyl optionally substituted with C_{1-4} alkyl;

R^3 is nitrile, carboxy, $-COOR^8$ where R^8 is an ester group, $-CONR^9R^{10}$ where R^9 and R^{10} are each hydrogen or C_{1-4} alkyl or $R^{11}SO_2$ where R^{11} is C_{1-4} alkyl or optionally substituted phenyl;


R^4 is $-NR^{12}R^{13}$, $-NHCOR^{12}$, $-N(COR^{12})_2$ or $-N=CHOCH_2R^{12}$ where R^{12} and R^{13} are each hydrogen or C_{1-4} alkyl optionally substituted with carboxy, or R^4 is



where X is C_{2-4} alkylene, or R^4 is $-NHSO_2R^{14}$ where R^{14} is C_{1-4} alkyl or optionally substituted phenyl; and

n is 0-2.

US5281619 discloses naphthopyrans for therapy of diabetic complications:



5 wherein,

R^1 is C_{1-4} alkoxy, OH or COOH;

R^2 is optionally substituted phenyl;


R^3 is nitrile, or R^3 is carboxy or $-COOR^8$ when R^2 is phenyl substituted with 3-nitro or 3-trifluoromethyl and R^8 is an ester group;

10 R^4 is $NR^{12}R^{13}$, $-NHCOR^{12}$, $-N(COR^{12})_2$ or $-N=CHOCH_2R^{12}$, wherein R^{12} and R^{13} are each H or C_{1-4} alkyl; and

n is 0-2.

EP599514 discloses the preparation of pyranoquinoline derivatives as inhibitors of cell proliferation:

15




wherein R^1 is optionally substituted phenyl or optionally substituted heteroaryl selected from thienyl, pyridyl, benzothienyl, quinolinyl, benzofuranyl or benzimidazolyl, or R^1 is furanyl optionally substituted with C_{1-4} alkyl;

20

R^2 is nitrile, carboxy, $-CO_2R^4$ wherein R^4 is an ester group, $-CON(R^5)R^6$ where R^5 and R^6 are independently H or C_{1-4} alkyl, or R^7SO_2 where R^7 is C_{1-4} alkyl or optionally substituted phenyl;

R^3 is $-NR^8R^9$, $-NHCOR^8$, $-N(CO_2R^8)_2$, $-N=CHOR^8$ where R^8 and R^9 are independently H or C_{1-4} alkyl, or $-NHSO_2R^{10}$ where R^{10} is C_{1-4} alkyl or optionally substituted phenyl, or


25



where X is C₂₋₄ alkylene; and

the ring P represents a pyridine fused to the benzopyran nucleus.

EP618206 discloses the preparation of naphthopyran and
 5 pyranoquinoline as immunosuppressants and cell proliferation inhibitors:



wherein,

A-B is CH₂CH₂ or CH=CH;

10 each R¹ is independently halo, carboxy, trifluoromethyl, hydroxy, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkylthio, hydroxy-C₁₋₄alkyl, hydroxy-C₁₋₄alkoxy, nitrogen-containing heterocyclyl, nitro, trifluoromethoxy, -COOR⁵ where R⁵ is an ester group, -COR⁶, -CONR⁶R⁷ or -NR⁶R⁷ where R⁶ and R⁷ are each hydrogen or C₁₋₄ alkyl;

15 R² is phenyl, naphthyl or heteroaryl selected from thienyl, pyridyl, benzothienyl, quinoliny, benzofuranyl or benzimidazolyl, wherein said phenyl, naphthyl and heteroaryl groups are optionally substituted, or R² is furanyl optionally substituted with C₁₋₄ alkyl;


20 R³ is nitrile, carboxy, -COOR⁸ where R⁸ is an ester group, -CONR⁹R¹⁰ where R⁹ and R¹⁰ are each hydrogen or C₁₋₄ alkyl, or -SO₂R¹¹ where R¹¹ is C₁₋₄ alkyl or optionally substituted phenyl-C₁₋₄ alkyl;

25 R⁴ is 1-pyrrolyl, 1-imidazolyl or 1-pyrazolyl, each of which is optionally substituted by one or two C₁₋₄ alkyl, carboxyl, hydroxyl-C₁₋₄alkyl or -CHO groups, or R⁴ is 1-(1,2,4-triazolyl), 1-(1,3,4-triazolyl) or 2-(1,2,3-triazolyl), each of which is optionally substituted by a C₁₋₄ alkyl or C₁₋₄ perfluoroalkyl group, or R⁴ is 1-tetrazolyl optionally substituted by C₁₋₄ alkyl;

X is a pyridine or a benzene ring; and

n is 0-2.

EP619314 discloses the preparation of 4-phenyl-4*H*-naphtho(2,1-*b*)-pyran derivatives:




wherein,

R₁ and R₂ are independently halo, trifluoromethyl, C₁-C₄ alkoxy, hydroxy, nitro, C₁-C₄ alkyl, C₁-C₄ alkylthio, hydroxy-C₁-C₄ alkyl, hydroxy-C₁-C₄alkoxy, trifluoromethoxy, carboxy, -COOR₈ where R₈ is an ester group, -COR₉, -CONR₉R₁₀ or -NR₉R₁₀ where R₉ and R₁₀ are each hydrogen or C₁-C₄ alkyl;

R₃ is nitrile, carboxy or -CO₂R₁₁ wherein R₁₁ is an ester group;


R₄ is -NR₁₂R₁₃, -NR₁₂COR₁₃, -N(COR₁₂)₂ or -N=CHOCH₂R₁₂ where R₁₂ and R₁₃ are each hydrogen or C₁₋₄ alkyl, or R₄ is




where X is C₂-C₄ alkylene, or R₄ is optionally substituted 1-pyrrolyl; and

m and n are each independently 0-2. The compounds are said to be useful for the treatment of restenosis, immune disease, and diabetic complications.

Smith, *et al.*, (*Bioorg. Med. Chem. Lett.* 5:2783-2788 (1995)) reported the anti-rheumatic potential of a series of 2,4-di-substituted-4*H*-naphtho[1,2-*b*]pyran-3-carbonitriles. They reported that 4-(3-nitrophenyl)-2-(*N*-succinimido)-4*H*-naphtho[1,2-*b*]pyran-3-carbonitrile has proved to be acid stable and still retains biological activity:



Birch, *et al.*, (*Diabetes* 45:642-650 (1996)) reported that LY290181, an inhibitor of diabetes-induced vascular dysfunction, blocks protein kinase C-stimulated transcriptional activation through inhibition of transcription factor binding to a phorbol response element.



LY290181

Panda, *et al.*, (*J. Biol. Chem.* 272: 7681-7687 (1997)) reported the suppression of microtubule dynamics by LY290181, which might be the potential mechanism for its antiproliferative action.

Wood, *et al.*, (*Mol. Pharmacol.* 52: 437-444 (1997)) reported that LY290181 inhibited mitosis and microtubule function through direct tubulin binding.

PCT published patent application WO9824427 disclosed antimicrotubule compositions and methods for treating or preventing inflammatory diseases. LY290181 was listed as an antimicrotubule agent.

SUMMARY OF THE INVENTION

The present invention is related to the discovery that 7,8-fused 4*H*-chromene and analogs, as represented in Formula I, are activators of the caspase cascade and inducers of apoptosis. Thus, an aspect of the present invention is directed to the use of compounds of Formula I as inducers of apoptosis.

A second aspect of the present invention is to provide a method for treating, preventing or ameliorating neoplasia and cancer by administering a compound of Formula I to a mammal in need of such treatment.

Many of the compounds within the scope of the present invention are novel compounds. Therefore, a third aspect of the present invention is to provide novel compounds of Formula I, and to also provide for the use of these novel compounds for treating, preventing or ameliorating neoplasia and cancer.


A fourth aspect of the present invention is to provide a pharmaceutical composition useful for treating disorders responsive to the induction of apoptosis, containing an effective amount of a compound of Formula I in admixture with one or more pharmaceutically acceptable carriers or diluents.

A fifth aspect of the present invention is directed to methods for the preparation of novel compounds of Formula I.

DETAILED DESCRIPTION OF THE INVENTION

The present invention arises out of the discovery that 7,8-fused 4*H*-chromene and analogs, as represented in Formula I, are potent and highly efficacious activators of the caspase cascade and inducers of apoptosis. Therefore, compounds of Formula I are useful for treating disorders responsive to induction of apoptosis.

Specifically, compounds useful in this aspect of the present invention are represented by Formula I:



or pharmaceutically acceptable salts or prodrugs thereof, wherein:

X is O, S or NR₆, wherein R₆ is hydrogen or optionally substituted alkyl;

Y is CN, COR₇, CO₂R₇ or CONR_xR_y, wherein R₇, R_x and R_y are
 5 independently hydrogen, C₁₋₁₀ alkyl, haloalkyl, aryl, fused aryl, carbocyclic, a
 heterocyclic group, a heteroaryl group, alkenyl, alkynyl, arylalkyl, arylalkenyl,
 arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl,
 carbocycloalkyl, heterocycloalkyl, hydroxyalkyl or aminoalkyl; or R_x and R_y
 are taken together with the nitrogen to which they are attached to form a
 10 heterocycle;

Z is NR₈R₉, NHCOR₈, N(COR₈)₂, N(COR₈)(COR₉), N=CHOR₈ or N=CHR₈,
 wherein R₈ and R₉ are independently H, C₁₋₄ alkyl or aryl, or R₈ and R₉ are
 combined together with the group attached to them to form a heterocycle;

R₁-R₂ are independently hydrogen, halo, haloalkyl, aryl, fused aryl,
 15 carbocyclic, a heterocyclic group, a heteroaryl group, C₁₋₁₀ alkyl, alkenyl,
 alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl,
 heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl,
 aminoalkyl, carboxyalkyl, nitro, amino, cyano, acylamido, hydroxy, thiol,
 acyloxy, azido, alkoxy, carboxy, methylenedioxy, carbonylamido or
 20 alkylthiol;


R₅ is hydrogen or C₁₋₁₀ alkyl;

A is optionally substituted and is aryl, heteroaryl, saturated carbocyclic,
 partially saturated carbocyclic, saturated heterocyclic, partially saturated
 heterocyclic, arylalkyl or heteroarylalkyl; and

25 B is optionally substituted and is a fused thiazole, oxazole, 2-imino-imidazole,
 2,1,3-thiadiazole-2-one, thiazole-2-one, oxazole-2-one, imidazole-2-thione, thiazole-
 2-thione, oxazole-2-thione, imidazoline, oxazoline, thiazoline, triazole, oxazine,
 oxazine-2,3-dione, or piperazine ring.

Preferred compounds of Formula I include compounds wherein A is
 30 phenyl, naphthyl, pyridyl, quinolyl, isoquinolyl, thienyl, furyl, pyrrolyl,

2-phenylethyl or cyclohexyl, any of which is optionally substituted. Other preferred compounds are wherein A is



wherein R₁₀-R₁₄ are independently hydrogen, halo, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, C₁₋₁₀ alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, aminoalkyl, carboxyalkyl, nitro, amino, cyano, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, methylenedioxy, carbonylamido or alkylthiol; or

R₁₀ and R₁₁, or R₁₁ and R₁₂, taken together with the atoms to which they are attached, form an aryl, heteroaryl, optionally substituted carbocyclic or optionally substituted heterocyclic group, wherein said group is optionally substituted.


For example, R₁₀ and R₁₁, or R₁₁ and R₁₂, may be taken together to form a structure selected from the group consisting of -OCH₂O-, -(CH₂)₃-, -(CH₂)₄-, -OCH₂CH₂O-, -CH₂N(R)CH₂-, -CH₂CH₂N(R)CH₂-, CH₂N(R)CH₂CH₂-, -CH=CH-CH=CH-, -N(R)-CH=CH-, -CH=CH-N(R)-, -O-CH=CH-, -CH=CH-O-, -S-CH=CH-, -CH=CH-S-, -N=CH-CH=CH-, -CH=N-CH=CH-, -CH=CH-N=CH-, -CH=CH-CH=N- and -N=CH-CH=N-, wherein R is hydrogen, C₁₋₁₀ alkyl, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl or aminoalkyl.

Preferably, R₅ is hydrogen. Preferably, X is O or S. Most preferably, X is O. Preferably, Z is NH₂. Preferably, Y is CN.

Optional substituents on B include alkyl, cycloalkylalkyl, hydroxyalkyl, epoxyalkyl, alkoxyalkyl, aminoalkyl and haloalkyl. Preferred

optional substituents on B include methyl, hydroxymethyl, cyclopropylmethyl, and 2-*N,N*-diethylaminoethyl. Preferably, B is *N*-methyl-oxazol-2-one.

A preferred embodiment is represented by Formula II:




or pharmaceutically acceptable salts or prodrugs thereof, wherein R_1 , R_2 , R_5 , X, Y, Z and A are as defined previously with respect to Formula I;

D, E, and F are $CR_{15}R_{16}$, C=O, C=S, C=NR₁₅, S=O, N, NR₁₅, O, or S, wherein R_{15} and R_{16} are independently hydrogen, alkyl or aryl, provided that the ring comprising D, E and F is not a furan, thiophene, pyrrole, dihydropyrrole, 1,3-dioxolane, cyclopentyl, imidazole or imidazol-2-one ring; and the dashed line represents a single bond or a double bond, with the proviso that both the dashed lines can not be a double bond at the same time.

Such compounds include substituted benzothiazopyrans, benzoxazolepyrans, benzo-(1,2-dihydro-imidazole)pyrans, benzo-(1,2-dihydro-thiazole)pyrans, benzo-(1,2-dihydro-oxazole)pyrans, benzo-(1,2-dihydro-2-oxo-imidazole)pyrans, benzo-(1,2-dihydro-2-oxo-thiazole)pyrans, benzo-2-oxo-oxazolepyrans, benzo-(1,2-dihydro-2-thioxo-imidazole)pyrans, benzo-(1,2-dihydro-2-imino-imidazole)pyrans, benzo-(1,3-dihydro-2-oxo-2,1,3-thiadiazole)pyrans, and benzotriazolepyrans.

Preferred compounds falling within the scope of Formula II include compounds wherein R_1 - R_2 are hydrogen. Preferably R_5 is hydrogen; X is O; Z is NH₂ and Y is CN. Preferably A is optionally substituted phenyl, naphthyl, pyridyl, quinolyl, isoquinolyl, thienyl, furyl, pyrrolyl, 2-phenylethyl or cyclohexyl.

Another preferred embodiment is represented by Formula III:



or pharmaceutically acceptable salts or prodrugs thereof, wherein R_1 , R_2 , R_5 , X , Y , Z and A are as defined previously with respect to Formula I; and D , E , F , and G are $CR_{15}R_{16}$, CO , N , O , or S , wherein R_{15} and R_{16} are defined above, provided that at least two of D , E , F , and G are hetero atoms and the ring comprising D , E , F and G is not a 1,4-dioxolane, pyrazine or 1,4-dihydro-2,3-dioxopyrazine ring; and the dashed lines represent a single bond or a double bond.

Such compounds include substituted benzoxazinepyrans and benzo-2,3-dioxo-oxazinepyrans.

Preferred compounds falling within the scope of Formula III include compounds wherein R_1 - R_2 are hydrogen. Preferably R_5 is hydrogen.

Exemplary preferred compounds that may be employed in the method of the invention include, without limitation:

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-thioxo-4,7,8,9-tetrahydro-imidazo[4,5-*h*]chromene;

2-Amino-3-cyano-4-(3-methoxyphenyl)-8-thioxo-4,7,8,9-tetrahydro-imidazo[4,5-*h*]chromene;

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-imino-4,7,8,9-tetrahydro-imidazo[4,5-*h*]chromene;

2-Amino-3-cyano-4-(3-methoxyphenyl)-8-imino-4,7,8,9-tetrahydro-imidazo[4,5-*h*]chromene;

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,7,9-trihydro-2,1,3-thiadiazole[4,5-*h*]chromene;

2-Amino-3-cyano-4-(3-methoxyphenyl)-8-oxo-4,7,9-trihydro-2,1,3-thiadiazole[4,5-*h*]chromene;

2-Amino-3-cyano-4-(3-methoxyphenyl)-8-oxo-4,7,8-trihydro-oxazo[4,5-*h*]chromene;

2-Amino-3-cyano-4-(3-methoxyphenyl)-4H-oxazo[4,5-*h*]chromene;

2-Amino-3-cyano-4-(3-methoxyphenyl)-4,7,8,9-tetrahydro-8,9-dioxo-oxazine[2,3-*h*]chromene;

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4,7,8,9-tetrahydro-8,9-dioxo-oxazine[2,3-*h*]chromene;

5 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-thiazo[4,5-*h*]chromene;

2-Amino-3-cyano-4-(3-methoxyphenyl)-4*H*-thiazo[4,5-*h*]chromene;

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-triazolo[4,5-*h*]chromene;

10 2-Amino-3-cyano-4-(3-methoxyphenyl)-4*H*-triazolo[4,5-*h*]chromene;

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,8,9-trihydro-oxazo[5,4-*h*]chromene;

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,8-dihydro-9-methyl-oxazo[5,4-*h*]chromene;

15 2-Methoxymethylimine-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-oxazo[5,4-*h*]chromene;

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,7,8-trihydro-oxazo[4,5-*h*]chromene;

20 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,8-dihydro-7-methyl-oxazo[4,5-*h*]chromene;

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-oxazo[4,5-*h*]chromene; and

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-oxazo[5,4-*h*]chromene.

25 The present invention is also directed to novel compounds within the scope of Formulae I-III. Exemplary preferred novel compounds of this invention include, without limitation:

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-thioxo-4,7,8,9-tetrahydro-imidazo[4,5-*h*]chromene;

30 2-Amino-3-cyano-4-(3-methoxyphenyl)-8-thioxo-4,7,8,9-tetrahydro-imidazo[4,5-*h*]chromene;

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-imino-4,7,8,9-tetrahydro-imidazo[4,5-*h*]chromene;

- 2-Amino-3-cyano-4-(3-methoxyphenyl)-8-imino-4,7,8,9-tetrahydro-imidazo[4,5-*h*]chromene;
- 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,7,9-trihydro-2,1,3-thiadiazole[4,5-*h*]chromene;
- 5 2-Amino-3-cyano-4-(3-methoxyphenyl)-8-oxo-4,7,9-trihydro-2,1,3-thiadiazole[4,5-*h*]chromene;
- 2-Amino-3-cyano-4-(3-methoxyphenyl)-8-oxo-4,7,8-trihydro-oxazole[4,5-*h*]chromene;
- 2-Amino-3-cyano-4-(3-methoxyphenyl)-4*H*-oxazole[4,5-*h*]chromene;
- 10 2-Amino-3-cyano-4-(3-methoxyphenyl)-4,7,8,9-tetrahydro-8,9-dioxo-oxazine[2,3-*h*]chromene;
- 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4,7,8,9-tetrahydro-8,9-dioxo-oxazine[2,3-*h*]chromene;
- 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-thiazole[4,5-*h*]chromene;
- 15 2-Amino-3-cyano-4-(3-methoxyphenyl)-4*H*-thiazole[4,5-*h*]chromene;
- 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-triazole[4,5-*h*]chromene;
- 2-Amino-3-cyano-4-(3-methoxyphenyl)-4*H*-triazole[4,5-*h*]chromene;
- 20 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,8,9-trihydro-oxazole[5,4-*h*]chromene;
- 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,8-dihydro-9-methyl-oxazole[5,4-*h*]chromene;
- 2-Methoxymethylimine-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-oxazole[5,4-*h*]chromene;
- 25 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,7,8-trihydro-oxazole[4,5-*h*]chromene;
- 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,8-dihydro-7-methyl-oxazole[4,5-*h*]chromene;
- 30 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-oxazole[4,5-*h*]chromene; and
- 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-oxazole[5,4-*h*]chromene.

Useful alkyl groups include straight-chained and branched C₁₋₁₀ alkyl groups, more preferably C₁₋₆ alkyl groups. Typical C₁₋₁₀ alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, *sec*-butyl, *tert*-butyl, 3-pentyl, hexyl and octyl groups, which can be optionally substituted.

5 Useful alkoxy groups include oxygen substituted by one of the C₁₋₁₀ alkyl groups mentioned above, which can be optionally substituted.

Useful alkylthio groups include sulphur substituted by one of the C₁₋₁₀ alkyl groups mentioned above, which can be optionally substituted. Also included are the sulfoxides and sulfones of such alkylthio groups.

10 Useful amino groups include -NH₂, -NHR₁₅ and -NR₁₅R₁₆, wherein R₁₅ and R₁₆ are C₁₋₁₀ alkyl or cycloalkyl groups, or R₁₅ and R₁₆ are combined with the N to form a ring structure, such as a piperidine, or R₁₅ and R₁₆ are combined with the N and other group to form a ring, such as a piperazine. The alkyl group can be optionally substituted.

15 Optional substituents on the alkyl groups include one or more halo, hydroxy, carboxyl, amino, nitro, cyano, C₁-C₆ acylamino, C₁-C₆ acyloxy, C₁-C₆ alkoxy, aryloxy, alkylthio, C₆-C₁₀ aryl, C₄-C₇ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₆-C₁₀ aryl(C₂-C₆)alkenyl, C₆-C₁₀ aryl(C₂-C₆)alkynyl, saturated and unsaturated heterocyclic or heteroaryl. Optional substituents on the aryl, aralkyl and heteroaryl groups include one or more halo, C₁-C₆ haloalkyl, C₆-C₁₀ aryl, C₄-C₇ cycloalkyl, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₆-C₁₀ aryl(C₁-C₆)alkyl, C₆-C₁₀ aryl(C₂-C₆)alkenyl, C₆-C₁₀ aryl(C₂-C₆)alkynyl, C₁-C₆ hydroxyalkyl, nitro, amino, ureido, cyano, C₁-C₆ acylamino, hydroxy, thiol, C₁-C₆ acyloxy, azido, C₁-C₆ alkoxy or carboxy.

25 Useful aryl groups include C₆₋₁₄ aryl, preferably C₆₋₁₀ aryl. Typical C₆₋₁₄ aryl groups include phenyl, naphthyl, phenanthrenyl, anthracenyl, indenyl, azulenyl, biphenyl, biphenylenyl and fluorenyl groups.

Useful cycloalkyl groups are C₃₋₈ cycloalkyl. Typical cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

30 Useful saturated or partially saturated carbocyclic groups are cycloalkyl groups as described above, as well as cycloalkenyl groups, such as cyclopentenyl, cycloheptenyl and cyclooctenyl.

Useful halo or halogen groups include fluorine, chlorine, bromine and iodine.

Useful arylalkyl groups include any of the above-mentioned C₁₋₁₀ alkyl groups substituted by any of the above-mentioned C₆₋₁₄ aryl groups. Preferably the arylalkyl group is benzyl, phenethyl or naphthylmethyl.

Useful haloalkyl groups include C₁₋₁₀ alkyl groups substituted by one or more fluorine, chlorine, bromine or iodine atoms, e.g., fluoromethyl, difluoromethyl, trifluoromethyl, pentafluoroethyl, 1,1-difluoroethyl, chloromethyl, chlorofluoromethyl and trichloromethyl groups.

Useful acylamino (acylamido) groups are any C₁₋₆ acyl (alkanoyl) attached to an amino nitrogen, e.g., acetamido, chloroacetamido, propionamido, butanoylamido, pentanoylamido and hexanoylamido, as well as aryl-substituted C₁₋₆ acylamino groups, e.g., benzoylamido, and pentafluoro-benzoylamido.

Useful acyloxy groups are any C₁₋₆ acyl (alkanoyl) attached to an oxy (–O–) group, e.g., formyloxy, acetoxy, propionoyloxy, butanoyloxy, pentanoyloxy and hexanoyloxy.

Useful saturated or partially saturated heterocyclic groups include tetrahydrofuranyl, pyranal, piperidinyl, piperazinyl, pyrrolidinyl, imidazolidinyl, imidazolyl, indolyl, isoindolyl, quinuclidinyl, morpholinyl, isochromanal, chromanyl, pyrazolidinyl, pyrazolyl, tetronoyl and tetramoyl groups.

Useful heteroaryl groups include thienyl, benzo[b]thienyl, naphtho[2,3-*b*]thienyl, thianthrenyl, furyl, pyranal, isobenzofuranyl, chromenyl, xanthenyl, phenoxanthiyl, 2*H*-pyrrolyl, pyrrolyl, oxazolyl, imidazolyl, pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indolizyl, isoindolyl, 3*H*-indolyl, indolyl, indazolyl, purinyl, 4*H*-quinolizyl, isoquinolyl, quinolyl, phthalzyl, naphthyridinyl, quinoxalinyl, cinnolyl, pteridinyl, carbazolyl, β -carbolyl, phenanthridinyl, acridinyl, perimidinyl, phenanthrolinyl, phenazinyl, isothiazolyl, phenothiazinyl, isoxazolyl, furazanyl, phenoxazinyl, 1,4-dihydroquinoxaline-2,3-dione, 7-aminoisocoumarin, pyrido[1,2-*a*]pyrimidin-4-one, 1,2-benzisoxazol-3-yl, benzimidazolyl, benzimidazol-2-one, benzoxazolyl, benzoxazol-2-one,

2-oxindolyl and 2-oxobenzimidazolyl. Where the heteroaryl group contains a nitrogen atom in a ring, such nitrogen atom may be in the form of an *N*-oxide, e.g., a pyridyl *N*-oxide, pyrazinyl *N*-oxide and pyrimidinyl *N*-oxide.


5 Certain of the compounds of the present invention may exist as stereoisomers including optical isomers. The invention includes all stereoisomers and both the racemic mixtures of such stereoisomers as well as the individual enantiomers that may be separated according to methods that are well known to those of ordinary skill in the art.

10 Examples of pharmaceutically acceptable addition salts include inorganic and organic acid addition salts, such as hydrochloride, hydrobromide, phosphate, sulphate, citrate, lactate, tartrate, maleate, fumarate, mandelate and oxalate; and inorganic and organic base addition salts with bases such as sodium hydroxy, Tris(hydroxymethyl)aminomethane (TRIS, tromethane) and *N*-methyl-glucamine.

15 Examples of prodrugs of the compounds of the invention include the simple esters of carboxylic acid containing compounds (e.g., those obtained by condensation with a C₁₋₄ alcohol according to methods known in the art); esters of hydroxy containing compounds (e.g., those obtained by condensation with a C₁₋₄ carboxylic acid, C₃₋₆ dioic acid or anhydride thereof, such as succinic and fumaric anhydrides according to methods known in the art);
20 imines of amino containing compounds (e.g., those obtained by condensation with a C₁₋₄ aldehyde or ketone according to methods known in the art); carbamate of amino containing compounds, such as those described by Leu, *et. al.*, (*J. Med. Chem.* 42:3623-3628 (1999)) and Greenwald, *et. al.*, (*J. Med. Chem.* 42:3657-3667 (1999)); and acetals and ketals of alcohol containing
25 compounds (e.g., those obtained by condensation with chloromethyl methyl ether or chloromethyl ethyl ether according to methods known in the art); and phosphonate and phosphono compounds (e.g., those obtained by condensation with a phosphate ester, phosphoryl chloride, or phosphoric acid), which
30 include pharmaceutically acceptable mono-basic and di-basic addition salts of the phosphono group, for example, organic bases such as amine bases, which include ammonia, piperidine and morpholine.

The compounds of this invention may be prepared using methods known to those skilled in the art, or the novel methods of this invention. In particular, the compounds of this invention with Formulae I-III can be prepared as illustrated by exemplary reaction in Scheme 1. Reaction of a 2,3-disubstituted phenol, such as 2,3-diaminophenol with a substituted benzaldehyde, such as 3-methoxybenzaldehyde and malononitrile in the presence a base, such as piperidine or *N,N*-diisopropylethylamine produced the 7,8-diamino chromene, which can then cyclize under different conditions to produce the 7,8-fused chromenes. For example, reaction of the 7,8-diamino chromene with 1,1'-carbonyldiimidazole (CDI) will produce the 2-oxo-imidazole-chromene compound.

Scheme 1




5

Compounds of this invention with Formulae I-III can be prepared as illustrated by exemplary reaction in Scheme 2. Reaction of the diamino-chromene with CS₂ in the presence of a base such as KOH produces the 2-thioxo-imidazo-chromene compound. Reaction of the diamino-chromene with cyanogen bromide (BrCN) produces the 2-imino-imidazo-chromene

compound. Reaction of the diamino-chromene with SOCl_2 in the presence of a base, such as pyridine produces the 2-oxo-thiadiazo-chromene compound.

Scheme 2




5

Compounds of this invention with Formulae I-III also can be prepared as illustrated by exemplary reaction in Scheme 3. Reaction of 3-amino-pyrocatechol with a substituted benzaldehyde such as 3-methoxybenzaldehyde and malononitrile in the presence a base such as piperidine or *N,N*-diisopropylethylamine produces the 7-amino-8-hydroxy chromene, which can then cyclized under different conditions to produce the 7,8-fused chromenes. For example, reaction of the 7-amino-8-hydroxy chromene with 1,1'-carbonyldiimidazole (CDI) produces the 2-oxo-oxazole-chromene compound. Reaction of the 7-amino-8-hydroxy chromene with oxalyl chloride produces the 2,3-dioxo-oxazine-chromene compound.

10

15

Scheme 3



5 An important aspect of the present invention is the discovery that compounds having Formulae I-III are activators of caspases and inducers of apoptosis. Therefore, these compounds are expected to be useful in a variety of clinical conditions in which there is uncontrolled cell growth and spread of abnormal cells, such as in the case of cancer.

10 Another important aspect of the present invention is the discovery that compounds having Formulae I-III are potent and highly efficacious activators of caspases and inducers of apoptosis in drug resistant cancer cells, such as breast and prostate cancer cells, which enables these compounds to kill these drug resistant cancer cells. In comparison, most standard anti-cancer drugs are not effective in killing drug resistant cancer cells under the same conditions.

15 Therefore, compounds of this invention are expected to be useful for the treatment of drug resistant cancer in animals.

The present invention includes a therapeutic method useful to modulate *in vivo* apoptosis or *in vivo* neoplastic disease, comprising administering to a subject in need of such treatment an effective amount of a compound, or a pharmaceutically acceptable salt or prodrug of the compound

20

of Formulae I-III, which functions as a caspase cascade activator and inducer of apoptosis.

5 The present invention also include a therapeutic method comprising administering to an animal an effective amount of a compound, or a pharmaceutically acceptable salt or prodrug of said compound of Formulae I-III, wherein said therapeutic method is useful to treat cancer, which is a group of diseases characterized by the uncontrolled growth and spread of abnormal cells. Such diseases include, but are not limited to, Hodgkin's disease, non-Hodgkin's lymphomas, acute and chronic lymphocytic leukemias, multiple
10 myeloma, neuroblastoma, breast carcinomas, ovarian carcinomas, lung carcinomas, Wilms' tumor, cervical carcinomas, testicular carcinomas, soft-tissue sarcomas, chronic lymphocytic leukemia, primary macroglobulinemia, bladder carcinomas, chronic granulocytic leukemia, primary brain carcinomas, malignant melanoma, small-cell lung carcinomas, stomach carcinomas, colon
15 carcinomas, malignant pancreatic insulinoma, malignant carcinoid carcinomas, malignant melanomas, choriocarcinomas, mycosis fungoides, head and neck carcinomas, osteogenic sarcoma, pancreatic carcinomas, acute granulocytic leukemia, hairy cell leukemia, neuroblastoma, rhabdomyosarcoma, Kaposi's sarcoma, genitourinary carcinomas, thyroid
20 carcinomas, esophageal carcinomas, malignant hypercalcemia, cervical hyperplasia, renal cell carcinomas, endometrial carcinomas, polycythemia vera, essential thrombocytosis, adrenal cortex carcinomas, skin cancer, and prostatic carcinomas.

25 In practicing the therapeutic methods, effective amounts of compositions containing therapeutically effective concentrations of the compounds formulated for oral, intravenous, local and topical application, for the treatment of neoplastic diseases and other diseases in which caspase cascade mediated physiological responses are implicated, are administered to an individual exhibiting the symptoms of one or more of these disorders. The
30 amounts are effective to ameliorate or eliminate one or more symptoms of the disorders. An effective amount of a compound for treating a particular disease is an amount that is sufficient to ameliorate, or in some manner reduce, the symptoms associated with the disease. Such amount may be administered as a

single dosage or may be administered according to a regimen, whereby it is effective. The amount may cure the disease but, typically, is administered in order to ameliorate the disease. Typically, repeated administration is required to achieve the desired amelioration of symptoms

5 In another embodiment, a pharmaceutical composition comprising a compound, or a pharmaceutically acceptable salt of said compound of Formulae I-III, which functions as a caspase cascade activator and inducer of apoptosis in combination with a pharmaceutically acceptable vehicle is provided.

10 Another embodiment of the present invention is directed to a composition effective to inhibit neoplasia comprising a compound, or a pharmaceutically acceptable salt or prodrug of said compound of Formulae I-III, which functions as a caspase cascade activator and inducer of apoptosis, in combination with at least one known cancer chemotherapeutic agent, or a
15 pharmaceutically acceptable salt of said agent. Examples of known anti-cancer agents, which can be used for combination therapy include, but not are limited to alkylating agents, such as busulfan, cis-platin, mitomycin C, and carboplatin; antimitotic agents, such as colchicine, vinblastine, paclitaxel, and docetaxel; topo I inhibitors, such as camptothecin and topotecan; topo II
20 inhibitors, such as doxorubicin and etoposide; RNA/DNA antimetabolites, such as 5-azacytidine, 5-fluorouracil and methotrexate; DNA antimetabolites, such as 5-fluoro-2'-deoxy-uridine, ara-C, hydroxyurea and thioguanine; and antibodies, such as Herceptin[®] and Rituxan[®]. Other known anti-cancer agents which can be used for combination therapy include melphalan, chlorambucil,
25 cyclophosphamide, ifosfamide, vincristine, mitoguazone, epirubicin, acliarubicin, bleomycin, mitoxantrone, elliptinium, fludarabine, octreotide, retinoic acid, tamoxifen and alanosine.

 In practicing the methods of the present invention, the compound of the invention may be administered together with at least one known
30 chemotherapeutic agent as part of a unitary pharmaceutical composition. Alternatively, the compound of the invention may be administered apart from the at least one known cancer chemotherapeutic agent. In one embodiment, the compound of the invention and the at least one known cancer

chemotherapeutic agent are administered substantially simultaneously, i.e. the compounds are administered at the same time or one after the other, so long as the compounds reach therapeutic levels in the blood at the same time. On another embodiment, the compound of the invention and the at least one known cancer chemotherapeutic agent are administered according to their individual dose schedule, so long as the compounds reach therapeutic levels in the blood.

Another embodiment of the present invention is directed to a composition effective to inhibit neoplasia comprising a bioconjugates of said compound of Formulae I-III, which functions as a caspase cascade activator and inducer of apoptosis, in bioconjugation with at least one known therapeutically useful antibodies, such as Herceptin[®] or Rituxan[®]; growth factors, such as DGF, NGF; cytokines, such as IL-2, IL-4; or any molecule that binds to cell surface. The antibodies and other molecules will deliver compound of Formulae I-III to its targets and make them effective anticancer agents. The bioconjugates also could enhance the anticancer effect of therapeutically useful antibodies, such as Herceptin[®] or Rituxan[®].

Similarly, another embodiment of the present invention is directed to a composition effective to inhibit neoplasia comprising a compound or a pharmaceutically acceptable salt or prodrug of said compound of Formulae I-III, which functions as a caspase cascade activator and inducer of apoptosis, in combination with radiation therapy. In this embodiment, the compound of the invention may be administered at the same time as the radiation therapy is administered or at a different time.

Yet another embodiment of the present invention is directed to a composition effective for post-surgical treatment of cancer, comprising a compound, or a pharmaceutically acceptable salt or prodrug of said compound of Formulae I-III, which functions as a caspase cascade activator and inducer of apoptosis. The invention also relates to a method of treating cancer by surgically removing the cancer and then treating the animal with one of the pharmaceutical compositions described herein.

A wide range of immune mechanisms operate rapidly following exposure to an infectious agent. Depending on the type of infection, rapid

clonal expansion of the T and B lymphocytes occurs to combat the infection. The elimination of the effector cells following an infection is one of the major mechanisms maintaining immune homeostasis. This deletion of reactive cells has been shown to be regulated by a phenomenon known as apoptosis.

5 Autoimmune diseases have been lately identified as a consequence of deregulated cell death. In certain autoimmune diseases, the immune system directs its powerful cytotoxic effector mechanisms against specialized cells, such as oligodendrocytes in multiple sclerosis, the beta cells of the pancreas in diabetes mellitus, and thyrocytes in Hashimoto's thyroiditis (Ohsako, S. & Elkon, K.B., *Cell Death Differ.* 6:13-21 (1999)). Mutations of the gene encoding the lymphocyte apoptosis receptor Fas/APO-1/CD95 are reported to be associated with defective lymphocyte apoptosis and autoimmune lymphoproliferative syndrome (ALPS), which is characterized by chronic, histologically benign splenomegaly and generalized lymphadenopathy,

10 hypergammaglobulinemia, and autoantibody formation. (Infante, A.J., *et al.*, *J. Pediatr.* 133:629-633 (1998) and Vaishnaw, A.K., *et al.*, *J. Clin. Invest.* 103:355-363 (1999)). It was reported that overexpression of Bcl-2, which is a member of the bcl-2 gene family of programmed cell death regulators with anti-apoptotic activity, in developing B cells of transgenic mice, in the presence of T cell dependent costimulatory signals, results in the generation of a modified B cell repertoire and in the production of pathogenic autoantibodies (Lopez-Hoyos, M., *et al.*, *Int. J. Mol. Med.* 1:475-483 (1998)). It is therefore evident that many types of autoimmune disease are caused by defects of the apoptotic process, and one treatment strategy would be to turn

20 on apoptosis in the lymphocytes that are causing autoimmune disease (O'Reilly, L.A. & Strasser, A., *Inflamm. Res.* 48:5-21 (1999)).

Fas-Fas ligand (FasL) interaction is known to be required for the maintenance of immune homeostasis. Experimental autoimmune thyroiditis (EAT), characterized by autoreactive T and B cell responses and a marked lymphocytic infiltration of the thyroid, is a good model to study the therapeutic effects of FasL. Batteux, F., *et al.*, (*J. Immunol.* 162:603-608 (1999)) reported that by direct injection of DNA expression vectors encoding FasL into the inflamed thyroid, the development of lymphocytic infiltration

30

of the thyroid was inhibited and induction of infiltrating T cells death was observed. These results show that FasL expression on thyrocytes may have a curative effect on ongoing EAT by inducing death of pathogenic autoreactive infiltrating T lymphocytes.

5 Bisindolylmaleimide VIII is known to potentiate Fas-mediated apoptosis in human astrocytoma 1321N1 cells and in Molt-4T cells, both of which were resistant to apoptosis induced by anti-Fas antibody in the absence of bisindolylmaleimide VIII. Potentiation of Fas-mediated apoptosis by bisindolylmaleimide VIII was reported to be selective for activated, rather
10 than non-activated, T cells, and was Fas-dependent. Zhou T., *et al.*, (*Nat. Med.* 5:42-48 (1999)) reported that administration of bisindolylmaleimide VIII to rats during autoantigen stimulation prevented the development of symptoms of T cell-mediated autoimmune diseases in two models, the Lewis rat model of experimental allergic encephalitis and the Lewis adjuvant arthritis model.
15 Therefore, the application of a Fas-dependent apoptosis enhancer, such as bisindolylmaleimide VIII may be therapeutically useful for the more effective elimination of detrimental cells and inhibition of T cell-mediated autoimmune diseases. Therefore, an effective amount of a compound, or a pharmaceutically acceptable salt or prodrug of the compound of Formulae I-
20 III, which functions as a caspase cascade activator and inducer of apoptosis, should be an effective treatment for autoimmune disease.

 Psoriasis is a chronic skin disease that is characterized by scaly red patches. Psoralen plus ultraviolet A (PUVA) is a widely used and effective treatment for psoriasis vulgaris and Coven, *et al.*, *Photodermatol.*
25 *Photoimmunol. Photomed.* 15:22-27 (1999), reported that lymphocytes treated with psoralen 8-MOP or TMP plus UVA displayed DNA degradation patterns typical of apoptotic cell death. Ozawa, *et al.*, *J. Exp. Med.* 189:711-718 (1999) reported that induction of T cell apoptosis could be the main mechanism by which 312-nm UVB resolves psoriasis skin lesions. Low doses
30 of methotrexate may be used to treat psoriasis to restore a clinically normal skin. Heenen, *et al.*, *Arch. Dermatol. Res.* 290:240-245 (1998), reported that low doses of methotrexate may induce apoptosis and this mode of action could explain the reduction in epidermal hyperplasia during treatment of psoriasis

with methotrexate. Therefore, an effective amount of a compound, or a pharmaceutically acceptable salt or prodrug of the compound of Formulae I-III, which functions as a caspase cascade activator and inducer of apoptosis, should be an effective treatment for hyperproliferative skin diseases, such as psoriasis.

Synovial cell hyperplasia is a characteristic of patients with rheumatoid arthritis (RA). Excessive proliferation of RA synovial cells, as well as defective in synovial cell death, might be responsible for the synovial cell hyperplasia. Wakisaka, *et al.*, *Clin. Exp. Immunol.* 114:119-128 (1998), found that although RA synovial cells could die via apoptosis through Fas/FasL pathway, apoptosis of synovial cells was inhibited by proinflammatory cytokines present within the synovium, and suggested that inhibition of apoptosis by the proinflammatory cytokines may contribute to the outgrowth of synovial cells, and lead to pannus formation and the destruction of joints in patients with RA. Therefore, an effective amount of a compound, or a pharmaceutically acceptable salt or prodrug of the compound of Formulae I-III, which functions as a caspase cascade activator and inducer of apoptosis, should be an effective treatment for rheumatoid arthritis.

There have been accumulation of convincing evidence that apoptosis plays a major role in promoting resolution of the acute inflammatory response. Neutrophils are constitutively programmed to undergo apoptosis, thus limiting their pro-inflammatory potential and leading to rapid, specific, and non-phlogistic recognition by macrophages and semi-professional phagocytes (Savill, J., *J. Leukoc. Biol.* 61:375-380 (1997)). Boirivant, *et al.*, *Gastroenterology* 116:557-565 (1999), reported that lamina propria T cells isolated from areas of inflammation in Crohn's disease, ulcerative colitis, and other inflammatory states manifest decreased CD2 pathway-induced apoptosis, and that studies of cells from inflamed Crohn's disease tissue indicate that this defect is accompanied by elevated Bcl-2 levels. Therefore, an effective amount of a compound, or a pharmaceutically acceptable salt or prodrug of the compound of Formulae I-III, which functions as a caspase cascade activator and inducer of apoptosis, should be an effective treatment for inflammation and inflammatory bowel disease.

Compositions within the scope of this invention include all compositions wherein the compounds of the present invention are contained in an amount that is effective to achieve its intended purpose. While individual needs vary, determination of optimal ranges of effective amounts of each component is within the skill of the art. Typically, the compounds may be administered to mammals, e.g., humans, orally at a dose of 0.0025 to 50 mg/kg, or an equivalent amount of the pharmaceutically acceptable salt thereof, per day, of the body weight of the mammal being treated for apoptosis-mediated disorders. Preferably, approximately 0.01 to approximately 10 mg/kg is orally administered to treat or prevent such disorders. For intramuscular injection, the dose is generally approximately one-half of the oral dose. For example, a suitable intramuscular dose would be approximately 0.0025 to approximately 25 mg/kg, and most preferably, from approximately 0.01 to approximately 5 mg/kg. If a known cancer chemotherapeutic agent is also administered, it is administered in an amount with is effective to achieve its intended purpose. The amounts of such known cancer chemotherapeutic agents effective for cancer are well known to those of skill in the art.

The unit oral dose may be comprised of approximately 0.01 to approximately 50 mg, preferably approximately 0.1 to approximately 10 mg of the compound of the invention. The unit dose may be administered one or more times daily as one or more tablets, each containing from approximately 0.1 to approximately 10, conveniently approximately 0.25 to 50 mg of the compound or its solvates.

In a topical formulation, the compound may be present at a concentration of approximately 0.01 to 100 mg per gram of carrier.

In addition to administering the compound as a raw chemical, the compounds of the invention may be administered as part of a pharmaceutical preparation containing suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries, which facilitate processing of the compounds into preparations that can be used pharmaceutically. Preferably, the preparations, particularly those preparations which can be administered orally and which can be used for the preferred type of administration, such as

tablets, dragees, and capsules, and also preparations which can be administered rectally, such as suppositories, as well as suitable solutions for administration by injection or orally, contain from approximately 0.01 to 99 percent, preferably from approximately 0.25 to 75 percent of active compound(s), together with the excipient.

Also included within the scope of the present invention are the non-toxic pharmaceutically acceptable salts of the compounds of the present invention. Acid addition salts are formed by mixing a solution of the particular apoptosis inducers of the present invention with a solution of a pharmaceutically acceptable non-toxic acid, such as hydrochloric acid, fumaric acid, maleic acid, succinic acid, acetic acid, citric acid, tartaric acid, carbonic acid, phosphoric acid, oxalic acid, and the like. Basic salts are formed by mixing a solution of the particular apoptosis inducers of the present invention with a solution of a pharmaceutically acceptable non-toxic base, such as sodium hydroxide, potassium hydroxide, choline hydroxide, sodium carbonate, Tris, *N*-methyl-glucamine, and the like.

The pharmaceutical compositions of the invention may be administered to any animal which may experience the beneficial effects of the compounds of the invention. Foremost among such animals are mammals, e.g., humans and veterinary animals, although the invention is not intended to be so limited.

The pharmaceutical compositions of the present invention may be administered by any means that achieve their intended purpose. For example, administration may be by parenteral, subcutaneous, intravenous, intramuscular, intraperitoneal, transdermal, buccal, intrathecal, intracranial, intranasal or topical routes. Administration, alternatively, or concurrently, may be by the oral route. The dosage administered will be dependent upon the age, health, and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment, and the nature of the effect desired.

The pharmaceutical preparations of the present invention are manufactured in a manner which is itself known, e.g., by means of conventional mixing, granulating, dragee-making, dissolving, or lyophilizing processes. Thus, pharmaceutical preparations for oral use can be obtained by

combining the active compounds with solid excipients, optionally grinding the resulting mixture and processing the mixture of granules, after adding suitable auxiliaries, if desired or necessary, to obtain tablets or dragee cores.

Suitable excipients are, in particular, fillers, such as saccharides, e.g.,
5 lactose or sucrose, mannitol or sorbitol; cellulose preparations and/or calcium phosphates, e.g., tricalcium phosphate or calcium hydrogen phosphate, as well as binders, such as starch paste, using, e.g., maize starch, wheat starch, rice starch, potato starch, gelatin, tragacanth, methyl cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose, and/or
10 polyvinyl pyrrolidone. If desired, disintegrating agents may be added such as the above-mentioned starches and also carboxymethyl-starch, cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof, such as sodium alginate. Auxiliaries are, above all, flow-regulating agents and lubricants, for example, silica, talc, stearic acid or salts thereof, such as magnesium stearate
15 or calcium stearate, and/or polyethylene glycol. Dragee cores are provided with suitable coatings which, if desired, are resistant to gastric juices. For this purpose, concentrated saccharide solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, polyethylene glycol and/or titanium dioxide, lacquer solutions and suitable organic solvents or solvent
20 mixtures. In order to produce coatings resistant to gastric juices, solutions of suitable cellulose preparations, such as acetylcellulose phthalate or hydroxypropylmethyl-cellulose phthalate, are used. Dye stuffs or pigments may be added to the tablets or dragee coatings, e.g., for identification or in order to characterize combinations of active compound doses.

25 Other pharmaceutical preparations, which can be used orally, include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active compounds in the form of granules, which may be mixed with fillers, such as lactose; binders, such as starches, and/or lubricants,
30 such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds are preferably dissolved or suspended in suitable liquids, such as fatty oils, or liquid paraffin. In addition, stabilizers may be added.

Possible pharmaceutical preparations, which can be used rectally include, e.g., suppositories, which consist of a combination of one or more of the active compounds with a suppository base. Suitable suppository bases are, e.g., natural or synthetic triglycerides, or paraffin hydrocarbons. In addition, it is also possible to use gelatin rectal capsules, which consist of a combination of the active compounds with a base. Possible base materials include, e.g., liquid triglycerides, polyethylene glycols, or paraffin hydrocarbons.

Suitable formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form, e.g., water-soluble salts and alkaline solutions. In addition, suspensions of the active compounds as appropriate oily injection suspensions may be administered. Suitable lipophilic solvents or vehicles include fatty oils, e.g., sesame oil, or synthetic fatty acid esters, e.g., ethyl oleate or triglycerides or polyethylene glycol-400 (the compounds are soluble in PEG-400) cremophor, or cyclodextrins. Aqueous injection suspensions may contain substances, which increase the viscosity of the suspension include, e.g., sodium carboxymethyl cellulose, sorbitol, and/or dextran. Optionally, the suspension may also contain stabilizers.

In accordance with one aspect of the present invention, compounds of the invention are employed in topical and parenteral formulations and are used for the treatment of skin cancer.

The topical compositions of this invention are formulated preferably as oils, creams, lotions, ointments and the like by choice of appropriate carriers. Suitable carriers include vegetable or mineral oils, white petrolatum (white soft paraffin), branched chain fats or oils, animal fats and high molecular weight alcohol (greater than C_{12}). The preferred carriers are those in which the active ingredient is soluble. Emulsifiers, stabilizers, humectants and antioxidants may also be included, as well as agents imparting color or fragrance, if desired. Additionally, transdermal penetration enhancers can be employed in these topical formulations. Examples of such enhancers can be found in U.S. Patent Nos. 3,989,816 and 4,444,762.

Creams are preferably formulated from a mixture of mineral oil, self-emulsifying beeswax and water in which mixture the active ingredient,

dissolved in a small amount of an oil such as almond oil, is admixed. A typical example of such a cream is one which includes approximately 40 parts water, approximately 20 parts beeswax, approximately 40 parts mineral oil and approximately 1 part almond oil.

Ointments may be formulated by mixing a solution of the active ingredient in a vegetable oil, such as almond oil with warm soft paraffin and allowing the mixture to cool. A typical example of such an ointment is one which includes approximately 30% almond oil and approximately 70% white soft paraffin by weight.

Having now fully described this invention, it will be understood by those of ordinary skill in the art that the same can be performed within a wide and equivalent range of conditions, formulations and other parameters without affecting the scope of the invention or any embodiment thereof. All patents, patent applications and publications cited herein are fully incorporated by reference herein in their entirety.

EXAMPLE 1

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*- triazolo[4,5-*h*]chromene

In a round bottle (20 mL) was placed 2,7,8-triamino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-chromene (320 mg, 0.8 mmol), glacial acetic acid (1.0 mL, 1.6 mmol) and water (1 mL). The mixture was slightly warmed and a clear solution was obtained. The bottle was placed in ice water and the contents cooled to approximately 5 °C, and a cold solution of sodium nitrite (55 mg, 0.8 mmol) in 0.5 mL H₂O was added. The mixture was stirred for approximately 3 h at room temperature and the color darkened. The mixture was extracted using ethyl acetate and the extract was evaporated. The residue was purified by column chromatography (silica gel, hexane:EtOAc, 2:1) to yield 147 mg (43 %) of the title compound. ¹H NMR (CD₃OD): 7.76 (d, *J* = 5.5, 1H), 7.16 (d, *J* = 2.4 Hz, 1H), 7.11 (d, *J* = 5.5 Hz, 1H), 7.03 (d, *J* = 2.4 Hz, 1H), 4.59 (s, 1H), 3.91 (s, 3H), 3.84 (s, 3H).

EXAMPLE 2

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,8,9-trihydro-oxazo[5,4-*h*]chromene

1,1'-Carbonyldiimidazole (17.1 mg, 0.11 mmol, 1.5 eq.) was added to a solution of 2-amino-3-cyano-7-hydroxy-8-amino-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-chromene (29.4 mg, 0.07 mmol) in toluene (1 mL). The reaction mixture was heated to approximately 60 °C and stirred for approximately 18 h. The solvent was evaporated and the crude solid was washed with methanol several times to remove the soluble imidazole. The desired compound was dried under vacuum to yield 14.7 mg of a white solid. ¹H NMR (CDCl₃): 7.03-6.98 (m, 2H), 6.91-6.86 (m, 1H), 6.81 (s, 2H), 6.78 (d, *J* = 0.7 Hz, 1H), 4.79 (s, 1H), 3.79 (s, 3H), 3.68 (s, 3H).

EXAMPLE 3

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,8-dihydro-9-methyl-oxazo[5,4-*h*]chromene

Cesium carbonate (14 mg, 0.04 mmol, 1 eq.) was added to a solution of 2-amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,8,9-trihydro-oxazo[5,4-*h*]chromene (19.7 mg, 0.04 mmol) in dimethylformamide (2 mL), followed by a solution of iodomethane (2M in *t*-butyl methyl ether) (22 µL, 0.04 mmol, 1 eq.). The reaction mixture was stirred at room temperature overnight. The solvent was evaporated and the crude material was purified by Biotage flash chromatography (cartridge 12S, SiO₂) (Dyax Corporation) using approximately 0 % to 2 % methanol in dichloromethane to yield 9.5 mg of the desired compound. ¹H NMR (CD₃OD): 7.00 (d, *J* = 8.4 Hz, 1H), 6.95 (d, *J* = 2.0 Hz, 1H), 6.91 (d, *J* = 2.0 Hz, 1H), 6.84 (m, 1H), 4.76 (s, 1H), 3.86 (s, 3H), 3.79 (s, 3H), 3.68 (s, 3H).

EXAMPLE 4

2-Methoxymethylimine-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-oxazo[5,4-*h*]chromene

5 *p*-Toluenesulfonic acid (3 mg) was added to a solution of 2-amino-3-cyano-7-hydroxy-8-amino-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-chromene (25.2 mg, 0.06 mmol) in toluene (1 mL) followed by methylorthoformate (1 mL). The reaction mixture was heated to approximately 50 °C and stirred for approximately 1 h. The mixture was poured into saturated aqueous sodium bicarbonate solution (10 mL). The aqueous layer was extracted with ethyl acetate (3 x 10 mL). The organic layers were combined, dried over sodium sulfate, filtered and concentrated. The crude product was purified by Biotage flash chromatography (cartridge 12S, SiO₂) using approximately 35 % ethyl acetate in hexane to yield 15.5 mg of the desired compound. ¹H NMR (CDCl₃): 8.59 (s, 1H), 8.07 (s, 1H), 7.27 (d, *J* = 8.5 Hz, 1H), 6.95 (d, *J* = 8.6 Hz, 1H), 6.84 (d, *J* = 2.1 Hz, 1H), 6.70 (d, *J* = 2.1 Hz, 1H), 4.85 (s, 1H), 3.93 (d, *J* = 0.7 Hz, 1H), 3.77 (s, 3H), 3.77 (s, 3H)

EXAMPLE 5

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,7,8-trihydro-oxazo[4,5-*h*]chromene

20 (a) 2,3-Dihydroxybenzoic acid: 2,3-Dimethoxybenzoic acid (2 g, 10.98 mmol) was dissolved in acetic acid (25 mL). Hydriodic acid (approximately 47 % solution in water) (25 mL) was added and the reaction mixture was heated under reflux overnight. The reaction was allowed to cool to room temperature and the solvent removed under vacuum. The yellow solid was dissolved in a minimum of water and extracted with ethyl acetate (4 x 50 mL). The organic layers were combined, washed with approximately 10% sodium thiosulfate solution (2 x 50 mL), water (50 mL) and brine (50 mL). The organic layer was dried over sodium sulfate, filtered and concentrated under vacuum to yield 1.68 g (99 %) of 2,3-dihydroxybenzoic acid as a white solid. ¹H NMR (CD₃OD): 7.34 (d, *J* = 8.0 Hz, 1H), 6.99 (d, *J* = 8.0 Hz, 1H), 6.72 (t, *J* = 8.0 Hz, 1H).

(b) 7-Hydroxy-3H-benzooxazol-2-one: Diphenylphosphorylazide (336 μ L, 1.56 mmol, 1.2 eq) was added to 2,3-dihydroxybenzoic acid (200 mg, 1.30 mmol) in pyridine (10 mL) and heated at approximately 75 °C for approximately 30 h. The reaction mixture was allowed to cool to room temperature and the reaction was quenched with water. Pyridine was removed under vacuum and the residue was partitioned between ethyl acetate (20 mL) and a solution of saturated aqueous sodium bicarbonate:water (1:1, 20 mL). The aqueous layer was extracted with ethyl acetate (3 x 20 mL). The organic layers were combined, washed with brine, dried over sodium sulfate, filtered and concentrated under vacuum to yield a yellow oil. The desired product was isolated by Biotage flash chromatography (cartridge 40S, SiO₂) using approximately 0 %, 15 %, 25 % and 50 % ethyl acetate in hexanes as eluant to yield 104 mg (53 %) of 7-hydroxy-3H-benzooxazol-2-one as a white powder. ¹H NMR (CD₃OD): 6.96 (dd, *J* = 8.4, 7.8 Hz, 1H), 6.59 (d, *J* = 8.4 Hz, 1H), 6.56 (d, *J* = 7.8 Hz, 1H).

(c) 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,7,8-trihydro-oxazo[4,5-*h*]chromene: 7-Hydroxy-3H-benzooxazol-2-one (300 mg, 1.98 mmol), 5-bromoveratraldehyde (486 mg, 1.98 mmol) and malononitrile (131 mg, 1.98 mmol) were dissolved in ethanol (12 mL). Piperidine (390 μ L, 3.96 mmol) was added and the reaction stirred at room temperature overnight. The precipitate was filtered and dried to yield 338.5 mg (38 %) of the desired 2-amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,7,8-trihydrooxazo[4,5-*h*]chromene as a light yellow powder. ¹H NMR (DMSO-*d*₆) 11.79 (s, 1H), 7.21 (brs, 2H), 6.98 (s, 1H), 6.91 (s, 1H), 6.85-6.79 (m, 2H), 4.79 (s, 1H), 3.78 (s, 3H), 3.68 (s, 3H).

EXAMPLE 6

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,8-dihydro-7-methyl-oxazo[4,5-*h*]chromene

2-amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,7,8-trihydro-oxazo[4,5-*h*]chromene (335 mg, 0.75 mmol) and cesium carbonate (270 mg, 0.83 mmol) were dissolved in anhydrous acetone (10 mL). Methyl

iodide (2M solution in *t*-BuOMe, 415 μ L, 0.83 mmol) was added and the reaction stirred at room temperature overnight. The solvent was removed under vacuum and the residue partitioned between water (20 mL) and ethyl acetate (60 mL). The aqueous layer was extracted with ethyl acetate (2 x 50 mL). The organic layers were combined, dried over sodium sulfate, filtered and concentrated to a yellow solid. The product was isolated by Biotage flash chromatography (cartridge 25S, SiO₂) using 0 to 1% methanol in dichloromethane to yield 292 mg (85 %) of the title compound as a light yellow solid. ¹H NMR (Acetone-d₆): 7.04-6.93 (m, 4H), 6.49 (brs, 2H), 4.83 (s, 1H), 3.86 (s, 3H), 3.76 (s, 3H), 3.39 (s, 3H).

EXAMPLE 7

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-oxazo[4,5-*h*]chromene

(a) 2,3-dihydroxyaniline: 2,3-dimethoxyaniline (500 mg, 3.26 mmol) was dissolved in acetic acid (10 mL). Hydriodic acid (approximately a 47 % solution in water) (10 mL) was added and the reaction was heated under reflux for approximately 8 h. The reaction was cooled to room temperature and stirred for approximately 3 days. The solvent was removed under vacuum and the yellow solid dissolved in water and neutralized with saturated aqueous sodium bicarbonate solution. The aqueous layer was extracted with ethyl acetate (4 x 20 mL). The organic layers were combined, washed with a solution of approximately 10 % aqueous sodium thiosulfate (30 mL), water (30 mL) brine (30 mL), dried over sodium sulfate, filtered and concentrated. The crude product was purified by Biotage (cartridge 40S, SiO₂) using approximately a 1 %, 2 % and 4% methanol solution in dichloromethane to yield 188 mg (46%) of 2,3-dihydroxyaniline as a beige solid. ¹H NMR (CD₃OD): 6.47 (t, *J* = 8.0 Hz, 1H), 6.28 (dd, *J* = 8.0, 1.6 Hz, 1H), 6.24 (dd, *J* = 8.0, 1.6 Hz, 1H).

(b) 2-Amino-3-cyano-7-amino-8-hydroxy-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-chromene: 2,3-dihydroxyaniline (80 mg, 0.64 mmol), 5-bromoveratraldehyde (157 mg, 0.64 mmol) and malononitrile (42 mg, 0.64 mmol) were dissolved in ethanol (4 mL). Piperidine (127 μ L, 1.28

mmol) was added and the reaction stirred at room temperature overnight. The reaction mixture was concentrated under vacuum and the desired product isolated by Biotage flash chromatography (cartridge 12M) with approximately 2 % and 5% methanol solutions in dichloromethane as eluant to yield 97 mg (36 %) of the desired product as a brown foamy solid. ¹H NMR (CD₃OD): 6.86 (s, 1H), 6.80 (s, 1H), 6.47 (d, *J* = 8.0 Hz, 1H), 6.29 (d, *J* = 8.0 Hz, 1H), 4.56 (s, 1H), 3.78 (s, 3H), 3.75 (s, 3H).

(c) 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-oxazo[4,5-*h*]chromene: 2-amino-3-cyano-7-amino-8-hydroxy-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-chromene (30 mg, 0.07 mmol) was dissolved in toluene (1 mL) and THF (1 mL). Trimethylorthoformate (24 μ L, 0.22 mmol, 3 eq), and *p*-toluenesulfonic acid (2.7 mg, 0.01 mmol, 0.2 eq) were added and the reaction stirred at approximately 50 °C overnight. The reaction mixture was allowed to cool to room temperature and quenched with saturated aqueous sodium bicarbonate solution (10 mL). The aqueous layer was extracted with ethyl acetate (3 x 20 mL). The organic layers were combined, dried over sodium sulfate, filtered and concentrated to yield a brown oil. The product was isolated by Biotage flash chromatography (cartridge 12S, SiO₂) using approximately 0 %, 2 % and 4 % methanol solutions in dichloromethane as eluant to yield 13.5 mg of a yellow oily solid. NMR showed impurities in the aliphatic region. The product was re-crystallized in a minimum of methanol. The crystals were washed with diethyl ether to yield 4.6 mg (15 %) of the title compound as a yellow solid. ¹H NMR (DMSO-*d*₆): 8.79 (s, 1H), 7.49 (d, *J* = 8.0 Hz, 1H), 7.28 (brs, 2H), 7.09 (d, *J* = 8.0 Hz, 1H), 7.02 (s, 1H), 6.95 (s, 1H), 4.93 (s, 1H), 3.79 (s, 3H), 3.68 (s, 3H).

EXAMPLE 8

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-oxazo[5,4-*h*]chromene

A similar procedure as Example 7c was used to prepare the title compound: ¹H NMR (acetone-*d*₆): 8.52 (d, *J* = 0.3 Hz, 1H), 7.42 (d, *J* = 8.5 Hz, 1H), 7.22-7.19 (m, 1H), 7.07-7.04 (m, 2H), 6.51 (brs, 2H), 4.93 (s, 1H), 3.85 (s, 3H), 3.76 (s, 3H).

EXAMPLE 9

Identification of Compounds as Caspase Cascade Activators and Inducers of Apoptosis in Solid Tumor Cells

Human breast cancer cell lines T-47D and ZR-75-1 were grown according to media component mixtures designated by American Type Culture Collection + 10 % FCS (Invitrogen Corporation), in approximately 5 % CO₂- 95 % humidity incubator at approximately 37 °C. T-47D and ZR-75-1 cells were maintained at a cell density between 30 and 80 % confluency at a cell density of 0.1 to 0.6 x 10⁶ cells/mL. Cells were harvested at approximately 600 x g and resuspended at approximately 0.65 x 10⁶ cells/mL into appropriate media plus approximately 10 % FCS. An aliquot of approximately 45 µl of cells was added to a well of a 96-well microtiter plate containing approximately 5 µl of a solution of approximately 10 % DMSO in RPMI-1640 media solution containing approximately 0.16 to 10 µM of the test compounds (0.016 to 1 µM final). An aliquot of approximately 45 µl of cells was added to a well of a 96-well microtiter plate containing approximately 5 µl of a solution of approximately 10 % DMSO in RPMI-1640 media solution without test compound as the control sample. The samples were mixed by agitation and then incubated at approximately 37 °C for approximately 24 h in approximately 5 % CO₂-95% humidity incubator. After incubation, the samples were removed from the incubator and approximately 50 µl of a solution containing approximately 20 µM of *N*-(Ac-DEVD)-*N'*-ethoxycarbonyl-R110 fluorogenic substrate, SEQ ID NO:1 (Cytovia, Inc.; U.S. Patent No. 6,335,429), comprising approximately 20 % sucrose (Sigma), 20 mM DTT (Sigma), 200 mM NaCl (Sigma), 40 mM Na PIPES buffer at approximate pH 7.2 (Sigma), and approximately 500 µg/ml lysolecithin (Calbiochem) was added. The samples were mixed by agitation and incubated at room temperature. Using a fluorescent plate reader (Model 1420 Wallac Instruments), an initial reading (T = 0) was made approximately 1-2 min after addition of the substrate solution, employing excitation at 485 nm and emission at 530 nm, to determine the background fluorescence of the control sample. After the 3 h incubation, the samples were read for fluorescence as above (T = 3 h).

Calculation:

The Relative Fluorescence Unit values (RFU) were used to calculate the sample readings as follows:

$$5 \quad \text{RFU}_{(T=3h)} - \text{Control RFU}_{(T=0)} = \text{Net RFU}_{(T=3h)}$$

10 The activity of caspase cascade activation was determined by the ratio of the net RFU value for the test compounds to that of control samples. The EC₅₀ (nM) was determined by a sigmoidal dose-response calculation (Prism 2.0, GraphPad Software Inc.). The caspase activity (Ratio) and potency (EC₅₀) are summarized in Table I:


TABLE I. CASPASE ACTIVITY AND POTENCY

Example #	T-47D		ZR-75-1	
	Ratio	EC50 (nM)	Ratio	EC50 (nM)
1	1.8	>1000	5.3	560
2	6.4	566.6	8.5	365.6
3	1.5	Inactive	2.7	2501.3
4	8.1	47.2	7.9	23.0
5	8.6	284.7	4.5	156.2
6	6.5	11.8		
7	7.8	68.7	9.8	43.0
8	7.9	35.1	7.2	12.9

15 Thus, these compounds are identified as potent caspase cascade activators and inducers of apoptosis in solid tumor cells.

WHAT IS CLAIMED IS:

1. A method of treating a disorder responsive to the induction of apoptosis in an animal suffering therefrom, comprising administering to a mammal in need of such treatment, an effective amount of a compound of Formula I:



or a pharmaceutically acceptable salt or prodrug thereof, wherein:

X is O, S or NR₆, wherein R₆ is hydrogen or optionally substituted alkyl;

Y is CN, COR₇, CO₂R₇ or CONR_xR_y, wherein R₇, R_x and R_y are independently hydrogen, C₁₋₁₀ alkyl, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl or aminoalkyl; or

R_x and R_y are taken together with the nitrogen to form a heterocycle;

Z is NR₈R₉, NHCOR₈, N(COR₈)₂, N(COR₈)(COR₉), N=CHOR₈ or N=CHR₈, wherein R₈ and R₉ are independently H, C₁₋₄ alkyl or aryl, or R₈ and R₉ are combined together with the group attached to them to form a heterocycle;


R₁-R₂ are independently hydrogen, halo, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, C₁₋₁₀ alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, aminoalkyl, carboxyalkyl, nitro, amino, cyano, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, methylenedioxy, carbonylamido or alkylthiol;

R₅ is hydrogen or C₁₋₁₀ alkyl;

A is optionally substituted and is aryl, heteroaryl, saturated carbocyclic, partially saturated carbocyclic, saturated heterocyclic, partially saturated heterocyclic or arylalkyl; and

5 B is optionally substituted and is a fused thiazole, oxazole, 2-imino-imidazole, 2,1,3-thiadiazol-2-one, thiazol-2-one, oxazol-2-one, oxazol-2-thione, imidazol-2-thione, thiazol-2-thione, imidazoline, oxazoline, thiazoline, triazole, oxazine, oxazine-2,3-dione or piperazine ring.

- 10 2. The method of claim 1, wherein B is *N*-methyloxazol-2-one.
3. The method of claim 1, wherein X is O.
4. The method of claim 1, wherein A is optionally substituted and selected from the group consisting of phenyl, naphthyl, quinolyl, isoquinolyl, 15 pyridyl, thienyl, furyl, pyrrolyl, 2-phenylethyl and cyclohexyl.
5. The method of claim 1, wherein X is O, Y is CN and Z is NH₂.
6. The method of claim 1, wherein R₅ is hydrogen.
- 20 7. The method of claim 1, wherein A is



and R₁₀-R₁₄ are independently hydrogen, halo, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, C₁₋₁₀ alkyl, alkenyl, 25 alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, aminoalkyl, carboxyalkyl, nitro, amino, cyano, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, methylenedioxy, carbonylamido or alkylthiol; or

R_{10} and R_{11} , or R_{11} and R_{12} , taken together with the atoms to which they are attached form an aryl, heteroaryl, optionally substituted carbocyclic or optionally substituted heterocyclic group, wherein said group is optionally substituted.

5

8. The method of claim 7, wherein R_{10} and R_{11} , or R_{11} and R_{12} , are taken together to form a structure selected from the group consisting of
 $-OCH_2O-$, $-(CH_2)_3-$, $-(CH_2)_4-$, $-OCH_2CH_2O-$, $-CH_2N(R)CH_2-$,
 $-CH_2CH_2N(R)CH_2-$, $-CH_2N(R)CH_2CH_2-$, $-CH=CH-CH=CH-$,
10 $-N(R)-CH=CH-$, $-CH=CH-N(R)-$, $-O-CH=CH-$, $-CH=CH-O-$,
 $-S-CH=CH-$, $-CH=CH-S-$, $-N=CH-CH=CH-$, $-CH=N-CH=CH-$,
 $-CH=CH-N=CH-$, $-CH=CH-CH=N-$ and $-N=CH-CH=N-$, wherein R is
hydrogen, C_{1-10} alkyl, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic
group, a heteroaryl group, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl,
15 heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl,
heterocycloalkyl, hydroxyalkyl or aminoalkyl.

9. The method of claim 8, wherein said compound is
2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-thioxo-4,7,8,9-
20 tetrahydro-imidazo[4,5-*h*]chromene;
2-Amino-3-cyano-4-(3-methoxyphenyl)-8-thioxo-4,7,8,9-tetrahydro-
imidazo[4,5-*h*]chromene;
2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-imino-4,7,8,9-
tetrahydro-imidazo[4,5-*h*]chromene;
25 2-Amino-3-cyano-4-(3-methoxyphenyl)-8-imino-4,7,8,9-tetrahydro-
imidazo[4,5-*h*]chromene;
2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,7,9-
trihydro-2,1,3-thiadiaz[4,5-*h*]chromene;
2-Amino-3-cyano-4-(3-methoxyphenyl)-8-oxo-4,7,9-trihydro-2,1,3-
30 thiadiaz[4,5-*h*]chromene;
2-Amino-3-cyano-4-(3-methoxyphenyl)-8-oxo-4,7,8-trihydro-
oxazo[4,5-*h*]chromene;
2-Amino-3-cyano-4-(3-methoxyphenyl)-4*H*-oxazo[4,5-*h*]chromene;

2-Amino-3-cyano-4-(3-methoxyphenyl)-4,7,8,9-tetrahydro-8,9-dioxo-oxazine[2,3-*h*]chromene;

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4,7,8,9-tetrahydro-8,9-dioxo-oxazine[2,3-*h*]chromene;

5 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-thiazo[4,5-*h*]chromene;

2-Amino-3-cyano-4-(3-methoxyphenyl)-4*H*-thiazo[4,5-*h*]chromene;

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-triazolo[4,5-*h*]chromene;

10 2-Amino-3-cyano-4-(3-methoxyphenyl)-4*H*-triazolo[4,5-*h*]chromene;

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,8,9-trihydro-oxazo[5,4-*h*]chromene;

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,8-dihydro-9-methyl-oxazo[5,4-*h*]chromene;

15 2-Methoxymethylimine-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-oxazo[5,4-*h*]chromene;

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,7,8-trihydro-oxazo[4,5-*h*]chromene;


20 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,8-dihydro-7-methyl-oxazo[4,5-*h*]chromene;

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-oxazo[4,5-*h*]chromene; and

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-oxazo[5,4-*h*]chromene.

25

10. The method of claim 1, wherein said compound has the Formula II:



30


or pharmaceutically acceptable salts or prodrugs thereof, wherein

D, E, and F are CR₁₅R₁₆, CO, N, O, or S, wherein R₁₅ and R₁₆, are independently hydrogen or alkyl; and

the dashed line represents a single bond or a double bond, with the proviso that both the dashed lines can not be a double bond at the same time.

5

11. The method of claim 1, wherein said compound has the formula:



10 or pharmaceutically acceptable salts or prodrugs thereof, wherein R₁, R₂, R₅, X, Y, Z and A are as defined previously with respect to Formula I; and

D, E, F, and G are CR₁₅R₁₆, CO, N, O, or S, wherein R₁₅ and R₁₆ are independently hydrogen or alkyl, provided that at least two of D, E, F, and G are hetero atoms;

15 the dashed lines represent a single bond or a double bond.

12. The method of claim 11, wherein said compound is

2-amino-3-cyano-4-(3-methoxyphenyl)-4,7,8,9-tetrahydro-8,9-dioxo-oxazine[2,3-*h*]chromene; or

20 2-amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4,7,8,9-tetrahydro-8,9-dioxo-oxazine[2,3-*h*]chromene.

13. The method of claim 1, wherein said disorder is cancer.

25 14. The method of claim 13, wherein said cancer is selected from the group consisting of Hodgkin's disease, non-Hodgkin's lymphoma, acute and chronic lymphocytic leukemias, multiple myeloma, neuroblastoma, breast carcinoma, ovarian carcinoma, lung carcinoma, Wilms' tumor, cervical carcinoma, testicular carcinoma, soft-tissue sarcoma, chronic lymphocytic

leukemia, primary macroglobulinemia, bladder carcinoma, chronic granulocytic leukemia, primary brain carcinoma, malignant melanoma, small-cell lung carcinoma, stomach carcinoma, colon carcinoma, malignant pancreatic insulinoma, malignant carcinoid carcinoma, malignant melanoma, choriocarcinoma, mycosis fungoides, head and neck carcinoma, osteogenic sarcoma, pancreatic carcinoma, acute granulocytic leukemia, hairy cell leukemia, neuroblastoma, rhabdomyosarcoma, Kaposi's sarcoma, genitourinary carcinoma, thyroid carcinoma, esophageal carcinoma, malignant hypercalcemia, cervical hyperplasia, renal cell carcinoma, endometrial carcinoma, polycythemia vera, essential thrombocytosis, adrenal cortex carcinoma, skin cancer and prostatic carcinoma.

15. The method of claim 13, wherein said cancer is a drug resistant cancer.

15

16. The method of claim 13, additionally comprising administering at least one known cancer chemotherapeutic agent, or a pharmaceutically acceptable salt of said agent.

17. The method of claim 16, wherein said known cancer therapeutic agent is selected from the group consisting of busulfan, cis-platin, mitomycin C, carboplatin, colchicine, vinblastine, paclitaxel, docetaxel, camptothecin, topotecan, doxorubicin, etoposide, 5-azacytidine, 5-fluorouracil, methotrexate, 5-fluoro-2'-deoxy-uridine, ara-C, hydroxyurea, thioguanine, melphalan, chlorambucil, cyclophosphamide, ifosfamide, vincristine, mitoguazone, epirubicin, aclarubicin, bleomycin, mitoxantrone, elliptinium, fludarabine, octreotide, retinoic acid, tamoxifen, Herceptin®, Rituxan® and alanosine.

18. The method of claim 13, additionally comprising treating with radiation-therapy.

19. The method of claim 13, wherein said compound is administered after surgical treatment for cancer.

20. The method of claim 1, wherein said disorder is an autoimmune disease.

21. The method of claim 1, wherein said disorder is rheumatoid arthritis.


22. The method of claim 1, wherein said disorder is inflammation.

23. The method of claim 1, wherein said disorder is inflammatory bowel disease.

24. The method of claim 1, wherein said disorder is psoriasis.

25. The method of claim 1, wherein said disorder is a skin disease.

26. A compound of Formula I:



or a pharmaceutically acceptable salt or prodrug thereof, wherein:

X is O, S or NR₆, wherein R₆ is hydrogen or optionally substituted alkyl;

Y is CN, COR₇, CO₂R₇ or CONR_xR_y, wherein R₇, R_x and R_y are independently hydrogen, C₁₋₁₀ alkyl, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl or aminoalkyl; or

R_x and R_y are taken together with the nitrogen to form a heterocycle;

Z is NR_8R_9 , $NHCOR_8$, $N(COR_8)_2$, $N(COR_8)(COR_9)$, $N=CHOR_8$ or $N=CHR_8$, wherein R_8 and R_9 are independently H, C_{1-4} alkyl or aryl, or R_8 and R_9 are combined together with the group attached to them to form a heterocycle;

R_1 - R_2 are independently hydrogen, halo, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, C_{1-10} alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, aminoalkyl, carboxyalkyl, nitro, amino, cyano, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, methylenedioxy, carbonylamido or alkylthiol;


R_5 is hydrogen or C_{1-10} alkyl;

A is optionally substituted and is aryl, heteroaryl, saturated carbocyclic, partially saturated carbocyclic, saturated heterocyclic, partially saturated heterocyclic or arylalkyl; and

B is optionally substituted and is a fused thiazole, oxazole, 2-imino-imidazole, 2,1,3-thiadiazol-2-one, thiazol-2-one, oxazol-2-one, oxazol-2-thione, imidazol-2-thione, thiazol-2-thione, imidazoline, oxazoline, thiazoline, triazole, oxazine, oxazine-2,3-dione, or piperazine ring.

27. The compound of claim 26, wherein B is *N*-methyloxazol-2-one.

28. A compound of Formula II:



or pharmaceutically acceptable salts or prodrugs thereof, wherein

X is O, S or NR_6 , wherein R_6 is hydrogen or optionally substituted alkyl;

Y is CN, COR₇, CO₂R₇ or CONR_xR_y, wherein R₇, R_x and R_y are independently hydrogen, C₁₋₁₀ alkyl, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl or aminoalkyl; or

R_x and R_y are taken together with the nitrogen to form a heterocycle;

Z is NR₈R₉, NHCOR₈, N(COR₈)₂, N(COR₈)(COR₉), N=CHOR₈ or N=CHR₈, wherein R₈ and R₉ are independently H, C₁₋₄ alkyl or aryl, or R₈ and R₉ are combined together with the group attached to them to form a heterocycle;

R₁-R₂ are independently hydrogen, halo, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, C₁₋₁₀ alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, aminoalkyl, carboxyalkyl, nitro, amino, cyano, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, methylenedioxy, carbonylamido or alkylthiol;

R₅ is hydrogen or C₁₋₁₀ alkyl;

A is optionally substituted and is aryl, heteroaryl, saturated carbocyclic, partially saturated carbocyclic, saturated heterocyclic, partially saturated heterocyclic or arylalkyl;

D, E and F are CR₁₅R₁₆, CO, N, O, or S, wherein R₁₅ and R₁₆ are independently hydrogen or alkyl, provided that the ring comprising D, E and F is not a furan, thiophene, pyrrole, dihydropyrrole, 1,3-dioxolane, cyclopentyl, imidazol-2-one or imidazole ring; and

the dashed line represents a single bond or a double bond, with the proviso that both the dashed lines can not be a double bond at the same time.


29. The compound of claim 28, wherein the ring comprising D, E and F is *N*-methyloxazol-2-one.

30. The compound of claim 28, wherein said compound is

- 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-thioxo-4,7,8,9-tetrahydro-imidazo[4,5-*h*]chromene;
- 2-Amino-3-cyano-4-(3-methoxyphenyl)-8-thioxo-4,7,8,9-tetrahydro-imidazo[4,5-*h*]chromene;
- 5 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-imino-4,7,8,9-tetrahydro-imidazo[4,5-*h*]chromene;
- 2-Amino-3-cyano-4-(3-methoxyphenyl)-8-imino-4,7,8,9-tetrahydro-imidazo[4,5-*h*]chromene;
- 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,7,9-trihydro-2,1,3-thiadiaz[4,5-*h*]chromene;
- 10 2-Amino-3-cyano-4-(3-methoxyphenyl)-8-oxo-4,7,9-trihydro-2,1,3-thiadiaz[4,5-*h*]chromene;
- 2-Amino-3-cyano-4-(3-methoxyphenyl)-8-oxo-4,7,8-trihydro-oxazo[4,5-*h*]chromene;
- 15 2-Amino-3-cyano-4-(3-methoxyphenyl)-4*H*-oxazo[4,5-*h*]chromene;
- 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-thiazo[4,5-*h*]chromene;
- 2-Amino-3-cyano-4-(3-methoxyphenyl)-4*H*-thiazo[4,5-*h*]chromene;
- 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-triaz[4,5-*h*]chromene;
- 20 triazo[4,5-*h*]chromene;
- 2-Amino-3-cyano-4-(3-methoxyphenyl)-4*H*-triaz[4,5-*h*]chromene;
- 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,8,9-trihydro-oxazo[5,4-*h*]chromene;
- 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,8-dihydro-9-methyl-oxazo[5,4-*h*]chromene;
- 25 2-Methoxymethylimine-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-oxazo[5,4-*h*]chromene;
- 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,7,8-trihydro-oxazo[4,5-*h*]chromene;
- 30 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-8-oxo-4,8-dihydro-7-methyl-oxazo[4,5-*h*]chromene;
- 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4*H*-oxazo[4,5-*h*]chromene; and

2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4H-oxazo[5,4-*h*]chromene.

31. A compound which has the Formula III:



or pharmaceutically acceptable salts or prodrugs thereof, wherein

X is O, S or NR₆, wherein R₆ is hydrogen or optionally substituted alkyl;

Y is CN, COR₇, CO₂R₇ or CONR_xR_y, wherein R₇, R_x and R_y are independently hydrogen, C₁₋₁₀ alkyl, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl or aminoalkyl; or

R_x and R_y are taken together with the nitrogen to form a heterocycle;

Z is NR₈R₉, NHCOR₈, N(COR₈)₂, N(COR₈)(COR₉), N=CHOR₈ or N=CHR₈, wherein R₈ and R₉ are independently H, C₁₋₄ alkyl or aryl, or R₈ and R₉ are combined together with the group attached to them to form a heterocycle;

R₁-R₂ are independently hydrogen, halo, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, C₁₋₁₀ alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, aminoalkyl, carboxyalkyl, nitro, amino, cyano, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, methylenedioxy, carbonylamido or alkylthiol;

R₅ is hydrogen or C₁₋₁₀ alkyl;

A is optionally substituted and is aryl, heteroaryl, saturated carbocyclic, partially saturated carbocyclic, saturated heterocyclic, partially saturated heterocyclic or arylalkyl;

D, E, F, and G are CR₁₅R₁₆, CO, N, O, or S, wherein CR₁₅ and R₁₆ are independently hydrogen or alkyl, provided that at least two of D, E, F, and G are hetero atoms and the ring comprising D, E, F and G is not a 1,4-dioxolane, 1,4-dihydro-2,3-dioxopyrazine or pyrazine ring; and

5 the dashed lines represent a single bond or a double bond.

32. The compound of claim 31, wherein said compound is

2-Amino-3-cyano-4-(3-methoxyphenyl)-4,7,8,9-tetrahydro-8,9-dioxo-oxazine[2,3-*h*]chromene; and

10 2-Amino-3-cyano-4-(3-bromo-4,5-dimethoxyphenyl)-4,7,8,9-tetrahydro-8,9-dioxo-oxazine[2,3-*h*]chromene.

33. A pharmaceutical composition, comprising the compound of any one of claims 26-32 and a pharmaceutically acceptable excipient or carrier.

15

34. The pharmaceutical composition of claim 33, further comprising at least one known cancer chemotherapeutic agent, or a pharmaceutically acceptable salt of said agent.

20

35. The pharmaceutical composition of claim 34, wherein said known cancer chemotherapeutic agent is selected from the group consisting of busulfan, cis-platin, mitomycin C, carboplatin, colchicine, vinblastine, paclitaxel, docetaxel, camptothecin, topotecan, doxorubicin, etoposide, 5-azacytidine, 5-fluorouracil, methotrexate, 5-fluoro-2'-deoxy-uridine, ara-C, hydroxyurea, thioguanine, melphalan, chlorambucil, cyclophosphamide, ifosfamide, vincristine, mitoguazone, epirubicin, aclarubicin, bleomycin, mitoxantrone, elliptinium, fludarabine, octreotide, retinoic acid, tamoxifen, Herceptin[®], Rituxan[®] and alanosine.

25

30

36. The pharmaceutical composition of claim 33, wherein said excipient or carrier is selected from the group consisting of saccharides, starch

pastes, gelatin, tragacanth, cellulose preparations, calcium phosphates and polyvinyl pyrrolidone.

-1-

SEQUENCE LISTING

<110> CYTOVIA, INC.
CAI, SUI XIONG
XU, LIFEN
STORER, RICHARD
ATTARDO, GIORGIO

<120> 7-8-FUSED 4H-CHROMENE AND ANALOGS AS ACTIVATORS OF CASPASES AND INDUCERS
OF APOPTOSIS AND THE USE THEREOF

<130> 1735.065PC01

<150> 60/290,976

<151> 2001-05-16

<160> 1

<170> PatentIn version 3.1

<210> 1

<211> 4

<212> PRT

<213> Artificial

<220>

<223> Fluorogenic substrate

<400> 1

Asp Glu Val Asp

1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/15398

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/4188, 31/4196, 31/424, 31/429, 31/436, 31/4365, 31/4375, 31/4985, 31/5365; C07D 471/04, 491/052, 487/04

US CL : 514/229.8, 250, 291, 292, 293, 359, 362, 366, 393; 544/101, 345; 546/82, 83; 548/135, 151, 218, 256, 302.1

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/229.8, 250, 291, 292, 293, 359, 362, 366, 393; 544/101, 345; 546/82, 83; 548/135, 151, 218, 256, 302.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
STN/CAS, structure search

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 618 206 A1 (LILLY INDUSTRIES LIMITED) 05 November 1994 (05.10.1994), whole document.	1-36

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"B" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&"

document member of the same patent family

Date of the actual completion of the international search

12 July 2002 (12.07.2002)

Date of mailing of the international search report

25 OCT 2002

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Richard L. Raymond

Telephone No. (703) 308-1235